

Meeting abstract

## Expression and activation of T cell receptor dependent transcription factors in regulatory T cells

H Bendfeldt\*, S Frischbutter, S Kroeger, A Radbruch and R Baumgrass

Address: Deutsches Rheumaforschungszentrum, Berlin, Germany

\* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

*Cell Communication and Signaling* 2009, **7**(Suppl 1):A1 doi:10.1186/1478-811X-7-S1-A1

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A1>

© 2009 Bendfeldt et al; licensee BioMed Central Ltd.

T cell receptor signalling and therefore production of IL-2 upon antigen stimulation, has been shown to be impaired in regulatory T (Treg) cells. Whether the expression and activation of the major transcription factors NFATc2, AP-1 and Nf- $\kappa$ B are affected too, has not yet been determined. We found a strikingly lower expression of all three factors in human Treg cells compared to memory Th cells, but their activation was unharmed. Interestingly, after stimulation with PMA/Ionomycin, thus bypassing upstream signalling events, we found a small Treg cell subset, that was able to overcome its anergic phenotype and produced IL-2. This subpopulation is characterized by higher NFATc2, AP-1 and Nf- $\kappa$ B and lower FOXP3 levels compared to IL-2 nonproducing Treg cells. Our Data suggests that IL-2 production in Treg cells is not switched off by genetic imprinting, but rather the amounts and ratios of the essential transcription factors NFATc2, AP-1, Nf- $\kappa$ B and FOXP3 are essential to prevent IL-2 production in Treg cells and thereby support their anergic phenotype despite a very strong stimulation.